


**ESD RECORD COPY**RETURN TO  
SCIENTIFIC & TECHNICAL INFORMATION DIVISION  
(ESTI), BUILDING 1211

COPY NR. \_\_\_\_\_ OF \_\_\_\_\_ COPIES

**SEQUENTIAL INFORMATION SEEKING: AN OPTIMAL STRATEGY AND OTHER RESULTS****TECHNICAL DOCUMENTARY REPORT NO. ESD-TDR-64-605****OCTOBER 1964****David M. Messick****ESTI PROCESSED**☒ **DDC TAB** ☐ **PROJ OFFICER**☐ **ACCESSION MASTER FILE**☐ \_\_\_\_\_**DATE** \_\_\_\_\_**ESTI CONTROL NR.** **AL** **46004****CY NR.** \_\_\_\_\_ **OF** \_\_\_\_\_ **CYS****DECISION SCIENCES LABORATORY  
ELECTRONIC SYSTEMS DIVISION  
AIR FORCE SYSTEMS COMMAND  
UNITED STATES AIR FORCE  
L. G. Hanscom Field, Bedford, Massachusetts****Project 4690, Task 469003****(Prepared under Contract No. AF 19 (628)-1610 by the Psychometric Laboratory, University of North Carolina, Chapel Hill, North Carolina.)****A00614685**

When US Government drawings, specifications or other data are used for any purpose other than a definitely related government procurement operation, the government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Donot return. Retain or destroy.

#### DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies from Defense Documentation Center (DDC). Orders will be expedited if placed through the librarian or other person designated to request documents from DDC.

Copies available at Office of Technical Services, Department of Commerce.


SEQUENTIAL INFORMATION SEEKING: AN OPTIMAL STRATEGY AND OTHER RESULTS

TECHNICAL DOCUMENTARY REPORT NO. ESD-TDR-64-605

OCTOBER 1964

David M. Messick

DECISION SCIENCES LABORATORY  
ELECTRONIC SYSTEMS DIVISION  
AIR FORCE SYSTEMS COMMAND  
UNITED STATES AIR FORCE  
L. G. Hanscom Field, Bedford, Massachusetts



Project 4690, Task 469003

(Prepared under Contract No. AF 19 (628)-1610 by the Psychometric Laboratory, University of North Carolina, Chapel Hill, North Carolina.)


1. The first of these is the  
fact that the system is  
not self-sufficient. It  
requires a constant  
input of energy from  
the outside world.  
This is a serious  
drawback.

2. The second is the  
fact that the system  
is not self-sufficient.  
It requires a constant  
input of energy from  
the outside world.

## FOREWORD

This study was conducted under Contract No. AF 19(628)-1610 at the Psychometric Laboratory, University of North Carolina, Chapel Hill, North Carolina. Dr Albert Amon served as principal investigator and Dr Anne Story, as contract monitor.

Work was performed under Project 4690 "Information Processing in Command and Control", Task 469003, "Human Information Processing Techniques".



## ABSTRACT

This paper presents an optimal strategy for sequential sampling from binomial distributions. The strategy presented is general in that it is a "multi-action" rather than a two-action procedure. While the major task is to estimate the proportion,  $p$ , of "successes" in a hypothetical, infinite population of binary observations, it is assumed that the decision maker is only concerned with which of a set of mutually exclusive and exhaustive subsets of the unit interval contains  $p$ . The derived strategy maximizes the decision-maker's gain without regard to error probabilities.

The important variable in determining a rule for ceasing to look at new data and making a decision is found to be the expected probability of being correct. The criterion involves only the economic aspects of the situation. A "no information" theorem is presented which shows that under some circumstances when a "success" or a "failure" on a given trial are equally probable, the probability of being correct after making the observation is identical to the probability of being correct before the observation was taken. Finally, an appealing derivation of the Beta-binomial probability function is given which suggests a more tractable computational procedure for the distribution and which illuminates its limiting distribution.

PUBLICATION REVIEW AND APPROVAL

This Technical Documentary Report has been reviewed and approved.

*for* *Ronald W. Quisler*  
HERBERT RUBENSTEIN  
Chief, Decision Techniques Division  
Decision Sciences Laboratory

*Roy Morgan*  
ROY MORGAN, Colonel, USAF  
Director  
Decision Sciences Laboratory

## SEQUENTIAL INFORMATION SEEKING: AN OPTIMAL STRATEGY AND OTHER RESULTS

### I. Introduction.

It is not uncommon for a person to be faced with the task of having to evaluate a body of information and to make a decision on the basis of his evaluation. A problem which naturally arises is how long to spend collecting or evaluating data before making the decision. This issue is particularly pertinent when data or information can be obtained by the decision maker but only at a cost. The decision maker must determine at what point he will cease looking at new data and make his decision. When a decision maker can at any time choose to observe more data or choose to make a decision, we say that the sampling procedure is sequential.

In the quest of psychologists to understand human behavior in general and human decision making in particular, the study of human information seeking is of prominent importance. The empirical research of Irwin and Smith (1956, 1957); Pruitt (1961); Lanzetta and Kanareff (1962); Edwards (1964); and Messick (1964a); has contributed to understanding of the problem.

Psychologists themselves are engaged in an almost continual process of collecting and evaluating data from empirical research. Wald (1947) has shown that sequential experiments are often more efficient in minimizing required sample size than experiments the size of which are predetermined. Fiske and Jones (1954) have emphasized this argument for psychologists, who as scientists, may benefit from sequential sampling procedures as research tools. For such purposes it is desired to find procedures which have specific properties, e.g., to minimize expected sample sizes holding error probabilities constant (Wald; 1947), to maximize expected utilities holding error probabilities constant (Edwards; 1964), or to reduce uncertainty to a preassigned level (DeGroot, 1962; Lindley, 1956, 1957).

While these two sources of interest in sequential sampling and information processing are distinct, the modern approach to the study of the "rational man" (called the "ideal observer" in contemporary psychophysics) is tending to bring the two interests together. Becker (1958), for example, studied human sequential sampling from the theoretical point of view of Wald. Edwards (1964) is studying the same behavior from a Bayesian position.

The primary purpose of the present report is to derive an optimal strategy for a special, but not uncommon situation, that of sequential sampling from binomial distributions. The strategy presented here is general in that it is a "multi-action" rather than a two-action procedure. Furthermore, while the major task is to estimate the proportion,  $p$ , of "successes" in an hypothetical, infinite population of binary observations, it is assumed that the decision maker is only concerned with which of a set of mutually exclusive and exhaustive subsets of the unit interval contains  $p$ . The derived strategy maximizes the decision-maker's gain without regard to error probabilities.

The results of this report have potential value as a general Bayesian research strategy. The immediate need to be filled by them is to provide a formal, rational model of information-seeking which can be used to evaluate the "optimality" of actual human information seeking behavior (Messick; 1964a, 1964b).

## II. An optimal Strategy for information seeking.

In deriving optimal strategies for information seeking, attention will be restricted to the case in which the prior probability distribution is of the Beta family, with density given by

$$(1) \quad f(p|\alpha, \beta) = B(\alpha, \beta)^{-1} p^{\alpha-1} (1-p)^{\beta-1}, \quad \alpha, \beta > 0,$$

and in which the data generating process is binomial. The restriction of the prior distribution to the Beta family has been adequately defended by Rapoport (1964) and Lindley (1957). The task of the decision maker is to select one of  $s$  mutually exclusive and exhaustive subsets of the unit interval which he believes contains  $p$ , the true proportion. We let this decision maker be motivated solely by the desire to maximize his expected gain and we assume that he uses Bayes formula to combine prior opinion with current information to produce a posteriori opinions. Thus, given a particular prior distribution with parameters  $(\alpha, \beta)$ , and given that our decision maker has observed  $r$  "successes" in  $n$  Bernoulli trials, then the state of his knowledge concerning  $p$  will be represented by a Beta distribution with parameters  $(\alpha+r, \beta+n-r)$ .

We will assume that the economic aspects of the situation have the following form: if the terminal decision is correct (i.e., if  $p$  is in the selected subset) the decision maker is given  $A$  dollars; if the terminal decision is incorrect he is fined  $L$  dollars; and each observation made costs  $C$  dollars. The expected gain of making a terminal decision after having observed  $r$  successes in  $n$  trials is

$$(2) \quad E(g|\alpha+r, \beta+n-r) = AP^*(\alpha+r, \beta+n-r) + L(1-P^*(\alpha+r, \beta+n-r)) + nc \\ = (A-L)P^*(\alpha+r, \beta+n-r) - L + nc,$$

where  $c$  is assumed negative,  $L < A$ , and where

$$(3) \quad P^*(\alpha+r, \beta+n-r) = \max_i \int_{I_i} f(p|\alpha+r, \beta+n-r) dp; \quad i = 1, 2, \dots, s.$$

$P^*(\alpha+r, \beta+n-r)$  is the maximum probability of being correct, where maximization is with respect to the set of terminal actions or decision categories,  $\{I_i\}$ .

To avoid notational difficulties let  $\alpha_n = \alpha+r$ ,  $\beta_n = \beta+n-r$ ,  $\gamma_n = \alpha_n + \beta_n = \alpha + \beta + n$ , and  $\delta_n = \alpha_n / \gamma_n$ . The question which must now be answered is what is the expected gain of taking another  $k$  observations and then making a terminal decision,  $k = 1, 2, \dots$ . Denote this expected gain  $E_k(g|\alpha_n, \beta_n)$ . It will be shown that

$$(4) \quad E_k(g|\alpha_n, \beta_n) = (A-L) \sum_{t=0}^k \Pr(t|k, \alpha_n, \beta_n) P^*(\alpha_n+t, \beta_n+k-t) + L + c(n+k).$$

In this formula  $\Pr(t|k, \alpha_n, \beta_n)$  gives the probability of obtaining  $t$  successes in the  $k$  trials when the prior Beta distribution has parameters  $(\alpha_n, \beta_n)$ . This probability is given by the Beta-binomial probability function (see Raiffa and Schlaifer, 1961, 237) defined by

$$(5) \quad \Pr(t|k, \alpha_n, \beta_n) = \int_0^1 b(t|k, p) f(p|\alpha_n, \beta_n) dp \\ = \frac{(t+\alpha_n-1)! (\beta_n+k-t-1)! k! (\gamma_n-1)!}{t!(k-t)! (\alpha_n-1)! (\beta_n-1)! (\gamma_n+k-1)!}.$$

(An intuitively appealing deviation of this probability function is given in Section IV.) Given that  $t$  successes have occurred in the  $k$  trials the expected gain is given by (2) and is found to be

$$(6) \quad E(g|\alpha_n+t, \beta_n+k-t) = (A-L) P^*(\alpha_n+t, \beta_n+k-t) + L + c(n+k).$$

The expression in (4) results from taking the expectation of (6) with respect to  $t$ .

In order to take a sample size that will maximize the expected gain, an additional observation is made if and only if there exists a  $k$  such that the difference

$$(7) \quad \Delta_k(\alpha_n, \beta_n) = E_k(g|\alpha_n, \beta_n) - E(g|\alpha_n, \beta_n) > 0, \quad k = 1, 2, 3, \dots$$

Letting

$$(8) \quad \Pi_k(\alpha_n, \beta_n) = \sum_{t=0}^k \Pr(t|k, \alpha_n, \beta_n) P^*(\alpha_n+t, \beta_n+k-t) - P^*(\alpha_n, \beta_n),$$

then

$$(9) \quad \Delta_k(\alpha_n, \beta_n) = (A-L)\Pi_k(\alpha_n, \beta_n) + ck.$$

Therefore a stopping rule equivalent to (7) is to take another observation if and only if for some  $k$

$$(10) \quad \Pi_k(\alpha_n, \beta_n) > -\frac{ck}{A-L}, \quad k = 1, 2, 3, \dots$$

Several interesting and useful properties of this sampling rule may be gleaned from (10). First, since  $\Pi_k(\alpha_n, \beta_n)$  is the difference between two probabilities it can range only between 1 and -1. Only positive values of  $\Pi_k$  would lead to additional sampling since the term on the right of the inequality will always be positive under our restrictions. Furthermore it is obvious that values of  $k$  greater than  $-\frac{A-L}{c}$  need not be considered since

$$k \geq -\frac{A-L}{c} \rightarrow -\frac{ck}{A-L} \geq 1.$$

Since  $\Pi_k$  is bounded above by 1, (10) will never hold for  $k \geq -\frac{A-L}{c}$ .

Feasible values of  $k$  can be further restricted by noting that  $P^*(\alpha_n, \beta_n) > 0$ . If we let  $1-P^*(\alpha_n, \beta_n) = \epsilon_n$ , then we note that (10) can be true only for values of  $k$  such that

$$(11) \quad k < -\epsilon_n \frac{(A-L)}{c}.$$

Thus only values of  $k$ ,  $k = 1, 2, 3, \dots, -\epsilon_n \frac{(A-L)}{c}$  need be tested.

Finally, as  $n$  increases without bound  $P^*(\alpha_n, \beta_n)$  approaches unity as a result of the fact that  $f(p|\alpha_n, \beta_n)$  approaches a point. Therefore,  $\epsilon_n$  goes to zero and

$$(12) \quad \lim_{n \rightarrow \infty} \epsilon_n \frac{(A-L)}{c} = 0.$$

Thus

$$(13) \quad \lim_{n \rightarrow \infty} \Pr(\text{sampling stops after } n \text{ observations}) = 1.$$

Exceptions to (13) occur only if  $(A-L)$  is infinitely large or if  $c = 0$ . These exceptions make good intuitive sense. If  $A$  is infinitely large or  $L$  infinitely negative, or if observations are free, then sampling might well be expected to continue for an indefinite period.

It will be of practical value to develop approximations to the rule given by (10). Two types of approximation will be suggested but research is needed to determine how good the approximations are. The difficulty with (10) is that in some cases it may be necessary to test  $\pi_k$  for many values of  $k$  before a decision can be made as to whether to take another observation. For example, in an experiment performed by Messick (1964)),  $A = 15$ ,  $L = 0$ , and  $c = .2$ . The maximum  $k$  is thus,  $\frac{A-L}{c} = 75$ . Clearly the procedure developed here would not be feasible if  $\pi_k$  had to be tested for all values of  $k$  between 1 and 75. One way in which this difficulty could be overcome is to test for  $k = 1, 2, 3, \dots, u$ , where  $u$  is predetermined by the experimenter on the basis of the degree of accuracy desired. An alternative procedure for approximating (10) would involve selecting more or less arbitrary values of  $k$ , perhaps by some random procedure. For example, one might test for  $k = 1, 2, 6, 16, 23$ . This latter procedure would be free of any bias which might be involved in testing only small values of  $k$ , but it would be more time consuming computationally.

### III. A "No Information" theorem.

Let  $I_1'$  be a subset of  $[0-1]$  such that  $I_1' = [x, 1-x]$ , and such that

$$(14) \quad P^*(\alpha, \beta) = \int_x^{1-x} f(p|\alpha, \beta) dp, \text{ when } \alpha = \beta; \text{ and}$$

$$(15) \quad P^*(\alpha+1, \beta) = \int_x^{1-x} f(p|\alpha+1, \beta) dp, \quad \text{and}$$

$$(16) \quad P^*(\alpha, \beta+1) = \int_x^{1-x} f(p|\alpha, \beta+1) dp.$$

Then,

$$(17) \quad P^*(\alpha, \beta) = P^*(\alpha+1, \beta) = P^*(\alpha, \beta+1) .$$

The equivalence of  $P^*(\alpha+1, \beta)$  and  $P^*(\alpha, \beta+1)$  follows from:

$$(18) \quad \int_0^x f(p|\alpha, \beta) dp = 1 - \int_0^{1-x} f(p|\beta, \alpha) dp.$$

The second equality in (17) will be assumed and a complete proof of the first will be given.

Writing (14) in full we have

$$P^*(\alpha, \beta) = P^*(\alpha, \beta) = \frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} \int_x^{1-x} p^{\alpha-1} (1-p)^{\alpha-1} dp.$$

However, since the Beta distribution is symmetric when  $\alpha=\beta$ , this can be written as

$$(19) \quad P^*(\alpha, \beta) = 1-2 \frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} \int_0^x p^{\alpha-1} (1-p)^{\alpha-1} dp .$$

Writing (15) in full we have

$$\begin{aligned}
 (20) \quad P^*(\alpha+1, \beta) &= \frac{\Gamma(2\alpha+1)}{\Gamma(\alpha+1)\Gamma(\alpha)} \int_x^{1-x} p^\alpha(1-p)^{\alpha-1} dp \\
 &= \frac{\Gamma(2\alpha+1)}{\Gamma(\alpha+1)\Gamma(\alpha)} \int_0^{1-x} p^\alpha(1-p)^{\alpha-1} dp - \frac{\Gamma(2\alpha+1)}{\Gamma(\alpha+1)\Gamma(\alpha)} \int_0^x p^\alpha(1-p)^{\alpha-1} dp \\
 &= 1 - \frac{\Gamma(2\alpha+1)}{\Gamma(\alpha)\Gamma(\alpha+1)} \int_0^x p^{\alpha-1}(1-p)^\alpha dp - \frac{\Gamma(2\alpha+1)}{\Gamma(\alpha+1)\Gamma(\alpha)} \int_0^x p^\alpha(1-p)^{\alpha-1} dp
 \end{aligned}$$

To prove (17) we show that (19) minus (20) is identically zero. First notice that

$$\frac{\Gamma(2\alpha+1)}{\Gamma(\alpha)\Gamma(\alpha+1)} = \frac{2\alpha\Gamma(2\alpha)}{\alpha\Gamma(\alpha)^2} = 2 \frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} .$$

Subtracting (20) from (19) we have

$$\begin{aligned}
 (21) \quad P^*(\alpha, \beta) - P^*(\alpha+1, \beta) &= -2 \frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} \int_0^x [p^{\alpha-1}(1-p)^{\alpha-1} - p^{\alpha-1}(1-p)^\alpha - p^\alpha(1-p)^{\alpha-1}] dp \\
 &= -2 \frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} \int_0^x (p^{\alpha-1}(1-p)^{\alpha-1})(1-(1-p) - p) dp \\
 &= 0.
 \end{aligned}$$

This theorem has the following interpretation. If  $\alpha=\beta$ , then the mean of the prior distribution is  $1/2$ . This value may be taken as the expected probability of a success on the next trial. If one's best terminal act at this point is to select the decision interval centered at  $1/2$ , and if this same terminal act remains optimal after another observation is taken, regardless of the outcome of the observation, then the probability of being correct before taking the observation is identical to the probability of being correct if the terminal act is selected after the one observation is made. From this point of view, the observation is non-informative.

#### IV. The Beta-binomial distribution: an intuitive derivation.

The binomial distribution,

$$(22) \quad b(r|n, p) = \binom{n}{r} p^r (1-p)^{n-r} ,$$

gives the probability of obtaining  $r$  "successes" in  $n$  independent Bernoulli trials when the probability of one success is  $p$ ,  $0 < p < 1$ . In this case,  $p$  is constant.

The Beta-binomial probability distribution,

$$(23) \quad \Pr(r|n, \alpha, \beta) = \int_0^1 b(r|n, p) f(p|\alpha, \beta) dp$$

$$= \frac{(r+\alpha-1)! (\beta+n-r-1)! n! (\alpha+\beta-1)!}{r!(n-r)! (\alpha-1)! (\beta-1)! (\alpha+\beta+n-1)!} ,$$

gives the probability of obtaining  $r$  successes in  $n$  trials when  $p$  is unknown, but rather treated as a random variable having a distribution of the Beta family with parameters  $(\alpha, \beta)$ .

To derive the Beta-binomial distribution, use will be made of the fact that the first moment of the Beta-distribution, which may be interpreted as the expected probability of a success, is given by

$$(24) \quad \delta = \frac{\alpha}{\alpha+\beta} ,$$

and therefore the expected probability of a "non-success" is

$$(25) \quad 1-\delta = \frac{\beta}{\alpha+\beta} .$$

If  $n$  observations are taken,  $r$  of which are successes, then the posterior Beta distribution has parameters  $(\alpha+r, \beta+n-r)$ . The mean of this posterior distribution is

$$(26) \quad \delta_n = \frac{\alpha+r}{\alpha+\beta+n} , \text{ and}$$

$$(27) \quad 1-\delta_n = \frac{\beta+n-r}{\alpha+\beta+n} .$$

We will only be concerned with the special case in which  $n = 1$ . Given  $(\alpha, \beta)$  we wish to find the probability of obtaining  $r$  successes in  $n$  trials.

To clarify the argument, consider the grid in Figure 1. A person begins at the origin with coordinates  $(\beta, \alpha)$ . If a success occurs our decision maker moves up one step. If a non-success occurs, he moves to the right one step. From any point on the grid, the probability of moving up a step is the ratio of the ordinate of the point to the sum of the coordinates and the probability of going to the right is the ratio of the abscissa to the sum of the coordinates.

We first note that the sum of the coordinates of any point is  $\alpha+\beta+n$ , where  $n$  is the number of steps (trials) required to reach the point from the origin. Second, the numerator of the probability of moving one step to the right does not depend on the ordinate and the numerator of the probability of moving one step up does not depend on the abscissa. Thus the numerator of the probability of moving from  $(\beta, \alpha)$  to  $(\beta+1, \alpha)$  is  $\beta$ , which is the same as the numerator of the probability of moving from  $(\beta, \alpha+4)$  to  $(\beta+1, \alpha+4)$ . Finally we note that the probability of going to any specified point in the grid from the origin via a specified path is the product of the probabilities of each of the component moves in the path.




Figure 1: Grid representing sequential sampling procedure. The origin is  $(\beta, \alpha)$ .

The major result to establish is that all paths to a point from the origin have the same probability. To prove this, first consider the numerator to the probability of moving from  $(\beta, \alpha)$  to  $(\beta+s, \alpha+r)$ . By the argument given in the above paragraph, the numerator of the probability of going  $s$  steps to the right and  $r$  steps up is given by  $\beta(\beta+1) \dots (\beta+s-1)\alpha(\alpha+1) \dots (\alpha+r-1)$  regardless of the order in which the steps occur. This number may be written as  $\frac{(\beta+s-1)!}{(\beta-1)!} \frac{(\alpha+r-1)!}{(\alpha-1)!}$ .

The denominator of the probability of moving in either direction from a point is the sum of the coordinates of the point, and, as stated above, this number depends only on the number of steps required to reach the point from the origin. Since the denominator of the product equals the product of the denominators of the step by step probabilities, it will be equal to  $(\alpha+\beta)(\alpha+\beta+1) \dots (\alpha+\beta+s+r-1)$ , which can be written  $\frac{(\alpha+\beta+s+r-1)!}{(\alpha+\beta-1)!}$ . Therefore, the probability of going from

$$(\alpha+\beta-1)!$$

$(\beta, \alpha)$  to  $(\beta+s, \alpha+r)$  by any specified path to

$$(28) \Pr[\text{any single path from } (\beta, \alpha) \text{ to } (\beta+s, \alpha+r)] =$$

$$\frac{(\beta+s-1)!}{(\beta-1)!} \frac{(\alpha+r-1)!}{(\alpha-1)!} \frac{(\alpha+\beta-1)!}{(\alpha+\beta+s+r-1)!}$$

Finally, the number of paths from  $(\beta, \alpha)$  to  $(\beta+s, \alpha+r)$  is

$$\binom{s+r}{r} = \binom{s+r}{s} = \frac{(s+r)!}{r! s!}$$

and the probability of going from  $(\beta, \alpha)$  to  $(\beta+s, \alpha+r)$  regardless of the path is

$$\Pr(r|r+s, \alpha, \beta) = \frac{(s+r)!}{r! s!} \frac{(\beta+s-1)!}{(\beta-1)!} \frac{(\alpha+r-1)!}{(\alpha-1)!} \frac{(\alpha+\beta-1)!}{(\alpha+\beta+s+r-1)!}$$

Setting  $n = s+r$  and  $s = n-r$ , the above expression is the Beta-binomial distribution given in (23).

This derivation of the Beta-binomial distribution is of practical value for two reasons. First, it provides a simpler computational procedure for finding probabilities than that in (23), which entails evaluating the factorials. Second, it becomes apparent from this point of view that the limiting distribution of the Beta-binomial is the binomial. As  $\alpha, \beta$  increase, the ratio  $\frac{\alpha}{\alpha+\beta}$  approaches  $p$  (see Raiffa and Schlaiffer, 1961). Therefore

$$(29) \lim_{\alpha, \beta \rightarrow \infty} \frac{\alpha(\alpha+1) \dots (\alpha+r-1) \beta(\beta+1) \dots (\beta+s-1)}{(\alpha+\beta)(\alpha+\beta+1) \dots (\alpha+\beta+s+r-1)} = p^r(1-p)^s = p^r(1-p)^{n-r}$$

#### Summary

In this report an optimal strategy is presented for sequential information seeking. The important variable in determining a stopping rule is found to be the expected probability of being correct. The criterion involves only the economic aspects of the situation. A "no information" theorem is presented which shows that under some circumstances when a "success" or a "failure" on a given trial are equally probable, the probability of being correct after making the observation is identical to the probability of being correct before the observation was taken. Finally, an appealing derivation of the Beta-binomial probability function was given which suggests a more tractable computational procedure for the distribution and which illuminates its limiting distribution.

### References

- Becker, G. M. Sequential decision making: Wald's model and estimates of parameters. J. exp. Psychol., 1958, 55, 629-636.
- DeGroot, M. H. Uncertainty, information, and sequential experiments. Ann. math. Statist., 1962, 33, 404-419.
- Edwards, W. Optimal Strategies for seeking information: models for statistics, choice reaction times, and human information processing. Institute of Science and Technology Report 3780-21-J, 1964, University of Michigan.
- Fiske, D. W. and Jones, L. V. Sequential analysis in psychological research. Psychol. Bull., 1954, 51, 264-275.
- Irwin, F. W. and Smith, W. A. S. Further tests of theories of decision in an "expanded judgement" situation. J. exp. Psychol., 1956, 52, 345-348.
- Irwin, F. W. and Smith, W. A. S. Value, cost, and information as determiners of decision. J. exp. Psychol., 1957, 54, 229-232.
- Lanzetta, J. T. and Kanareff, V. T. Information cost, amount of payoff, and level of aspiration as determinants of information seeking in decision making. Behav. Sci., 1962, 7, 459-473.
- Lindley, D. V. On a measure of the information provided by an experiment. Ann. math. Statist., 1956, 27, 986-1005.
- Lindley, D. V. Binomial sampling schemes and the concept of information. Biometrika, 1957, 44, 179-186.
- Messick, D. M. Sequential information seeking: effects of the number of terminal acts and prior information. Psychometric Lab. Rept. No. 41, University of North Carolina at Chapel Hill, 1964a.
- Messick, D. M. Information seeking in a computer controlled task: instructions, computer program, and data. Psychometric Lab. Research Memo. No. 18, University of North Carolina at Chapel Hill, 1964b.
- Pruitt, D. G. Informational requirements in making decisions. Amer. J. Psychol., 1961, 74, 433-439.
- Raiffa, H. and Schlaifer, R. Applied statistical decision theory. Boston: Graduate School of Business Administration, Harvard University, 1961.
- Rapoport, A. Sequential decision making in a computer-controlled task. J. math. Psychol., 1964, in press.
- Wald, A. Sequential Analysis. New York: John Wiley, 1947.



Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Psychometric Laboratory, University of North Carolina Chapel Hill, North Carolina		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP n/a	
3. REPORT TITLE  SEQUENTIAL INFORMATION SEEKING: AN OPTIMAL STRATEGY AND OTHER RESULTS			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) none			
5. AUTHOR(S) (Last name, first name, initial)  Messick, David M.			
6. REPORT DATE October 1964		7a. TOTAL NO. OF PAGES 12	7b. NO. OF REFS 15
8a. CONTRACT OR GRANT NO. AF 19(628)-1610		8a. ORIGINATOR'S REPORT NUMBER(S)  42	
b. PROJECT NO. 4690			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		ESD-TDR-64-605	
10. AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain from DDC OTS release authorized			
11. SUPPLEMENTARY NOTES  none		12. SPONSORING MILITARY ACTIVITY Decision Sciences Laboratory, Electronic Systems Division, L.G. Hanscom Field, Bedford, Massachusetts	
13. ABSTRACT <p>This paper presents an optimal strategy for sequential sampling from binomial distributions. The strategy presented is general in that it is a "multi-action" rather than a two-action procedure. While the major task is to estimate the proportion, <math>p</math>, of "successes" in a hypothetical, infinite population of binary observations, it is assumed that the decision maker is only concerned with which of a set of mutually exclusive and exhaustive subsets of the unit interval contains <math>p</math>. The derived strategy maximizes the decision-maker's gain without regard to error probabilities.</p> <p>The important variable in determining a rule for ceasing to look at new data and making a decision is found to be the expected probability of being correct. The criterion involves only the economic aspects of the situation. A "no information" theorem is presented which shows that under some circumstances when a "success" or a "failure" on a given trial are equally probable, the probability of being correct after making the observation is identical to the probability of being correct before the observation was taken. Finally, an appealing derivation of the Beta-binomial probability function is given which suggests a more tractable computational procedure for the distribution and which illuminates its limiting distribution.</p>			

DD FORM 1473  
1 JAN 64

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Decision Theory						

#### INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

